

AUTOMORPHISMS OF THE CATEGORY OF THE FREE NILPOTENT GROUPS OF THE FIXED CLASS OF NILPOTENCY.

A.Tsurkov

*Department of Mathematics and Statistics,
Bar Ilan University,
Ramat Gan, 52900, Israel.*

*Jerusalem College of Technology,
21 Havaad Haleumi,
Jerusalem, 91160, Israel.
tsurkov@jct.ac.il*

February 2, 2008

Abstract

This research was motivated by universal algebraic geometry. One of the central questions of universal algebraic geometry is: when two algebras have the same algebraic geometry? For answer of this question (see [8],[10]) we must consider the variety Θ , to which our algebras belongs, the category Θ^0 of all finitely generated free algebras of Θ and research how the group $\text{Aut}\Theta^0$ of all the automorphisms of the category Θ^0 are different from the group $\text{Inn}\Theta^0$ of the all inner automorphisms of the category Θ^0 . An automorphism Υ of the arbitrary category $\mathfrak{K}$ is called inner, if it is isomorphic as functor to the identity automorphism of the category $\mathfrak{K}$, or, in details, for every $A \in \text{Ob}\mathfrak{K}$ there exists $s_A^\Upsilon : A \rightarrow \Upsilon(A)$ isomorphism of these objects of the category $\mathfrak{K}$ and for every $\alpha \in \text{Mor}_\mathfrak{K}(A, B)$ the diagram

$$\begin{array}{ccc} A & \xrightarrow{s_A^\Upsilon} & \Upsilon(A) \\ \downarrow \alpha & & \Upsilon(\alpha) \downarrow \\ B & \xrightarrow{s_B^\Upsilon} & \Upsilon(B) \end{array}$$

is commutative. In the case when Θ is a variety of all groups we have the classical results which let us resolve this problem by an indirect way. In [3] proved that for every free group F_i the group $\text{Aut}(\text{Aut}F_i)$ coincides with the group $\text{Inn}(\text{Aut}F_i)$, from this result in [4] was concluded that $\text{Aut}(\text{End}F_i) = \text{Inn}(\text{End}F_i)$ and from this fact by theorem of reduction [2] it can be concluded that $\text{Aut}\Theta^0 = \text{Inn}\Theta^0$. In the case when $\Theta = \mathfrak{N}_d$

is a variety of the all nilpotent class no more then d groups we know by [6], that if the number of generators i of the free nilpotent class d group NF_i^d are bigger enough than d , then $\text{Aut}(\text{Aut}NF_i^d) = \text{Inn}(\text{Aut}NF_i^d)$. But we have no description of $\text{Aut}(\text{End}NF_i^d)$ and so can not use the theorem of reduction. In this paper the method of verbal operations is used. This method was established in [9]. In [9] by this method was very easily proved that $\text{Aut}\Theta^0 = \text{Inn}\Theta^0$ when Θ is the variety of all groups and the variety of all abelian groups. In this paper we will prove, by this method, that $\text{Aut}\mathfrak{N}_d^0 = \text{Inn}\mathfrak{N}_d^0$ for every $d \geq 2$.

1 Introduction and methodology.

This research was motivated by universal algebraic geometry. One of the central questions of universal algebraic geometry is: when two algebras have the same algebraic geometry? For answer of this question (see [8],[10]) we must consider the variety Θ , to which our algebras belongs, the category Θ^0 of all finitely generated free algebras of Θ and research how the group $\text{Aut}\Theta^0$ of all the automorphisms of the category Θ^0 are different from the group $\text{Inn}\Theta^0$ of the all inner automorphisms of the category Θ^0 . An automorphism Υ of the arbitrary category $\mathfrak{K}$ is called inner, if it is isomorphic as functor to the identity automorphism of the category $\mathfrak{K}$, or, in details, for every $A \in \text{Ob}\mathfrak{K}$ there exists $s_A^\Upsilon : A \rightarrow \Upsilon(A)$ isomorphism of these objects of the category $\mathfrak{K}$ and for every $\alpha \in \text{Mor}_\mathfrak{K}(A, B)$ the diagram

$$\begin{array}{ccc} A & \xrightarrow{s_A^\Upsilon} & \Upsilon(A) \\ \downarrow \alpha & & \Upsilon(\alpha) \downarrow \\ B & \xrightarrow{s_B^\Upsilon} & \Upsilon(B) \end{array}$$

is commutative. In the case when Θ is a variety of all groups we have the classical results which let us resolve this problem by an indirect way. In [3] proved that for every free group F_i the group $\text{Aut}(\text{Aut}F_i)$ coincides with the group $\text{Inn}(\text{Aut}F_i)$, from this result in [4] was concluded that $\text{Aut}(\text{End}F_i) = \text{Inn}(\text{End}F_i)$ and from this fact by theorem of reduction [2] it can be concluded that $\text{Aut}\Theta^0 = \text{Inn}\Theta^0$. In the case when $\Theta = \mathfrak{N}_d$ is a variety of the all nilpotent class no more then d groups we know by [6], that if the number of generators i of the free nilpotent class d group NF_i^d are bigger enough than d , then $\text{Aut}(\text{Aut}NF_i^d) = \text{Inn}(\text{Aut}NF_i^d)$. But we have no description of $\text{Aut}(\text{End}NF_i^d)$ and so can not use the theorem of reduction. In this paper the method of verbal operations is used. This method was established in [9]. In [9] by this method was very easily proved that $\text{Aut}\Theta^0 = \text{Inn}\Theta^0$ when Θ is the variety of all groups and the variety of all abelian groups. In this paper we will prove, by this method, that $\text{Aut}\mathfrak{N}_d^0 = \text{Inn}\mathfrak{N}_d^0$ for every $d \geq 2$.

We start the explanation of the method of verbal operations in the general situation. We consider the variety Θ of one-sorted algebras. The signature of our algebras we denote Ω .

For the construction of the category Θ^0 we must fix a countable set of symbols $X_0 = \{x_1, x_2, \dots, x_n, \dots\}$. As objects of the category Θ^0 we consider all free algebras $W(X)$ of the variety Θ generated by the finite subsets $X \subset X_0$. Morphisms of the category Θ^0 are homomorphisms of these algebras.

Definition 1.1. *We call the automorphism Φ of the category Θ^0 **strongly stable** if it fulfills these three conditions:*

A1) Φ preserves all objects of Θ^0 ,

A2) there exists a system of bijections $\{s_B^\Phi : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$ such that Φ acts on the morphisms of Θ^0 by these bijections, i. e.,

$$\Phi(\alpha) = s_B^\Phi \alpha (s_A^\Phi)^{-1} \quad (1.1)$$

for every $\alpha : A \rightarrow B$ ($A, B \in \text{Ob}\Theta^0$);

A3) $s_B^\Phi|_X = \text{id}_X$ for every $B = W(X) \in \text{Ob}\Theta^0$.

The variety Θ is called an IBN variety if for every free algebras $W(X), W(Y) \in \Theta$ we have $W(X) \cong W(Y)$ if and only if $|X| = |Y|$. In the [9, Theorem 2] proved, that if Θ is an IBN variety of one-sorted algebras, then every automorphism $\Psi \in \text{Aut}\Theta^0$ can be decomposed: $\Psi = \Upsilon\Phi$, where $\Upsilon, \Phi \in \text{Aut}\Theta^0$, Υ is an inner automorphism and Φ is a strongly stable one. So, if we want to know the difference of the group $\text{Aut}\Theta^0$ from the group $\text{Inn}\Theta^0$, we must study the strongly stable automorphisms of the category Θ^0 .

Before the explanation of the notion of the verbal operation we will introduce the short notation, which will be widely used in this paper. In this notation k -tuple $(c_1, \dots, c_k) \in C^k$ (C is an arbitrary set) we denote by single letter c and we will even allow ourself to write $c \in C$ instead $c \in C^k$ and to write "homomorphism $\alpha : A \ni a \rightarrow b \in B$ " instead "homomorphism $\alpha : A \rightarrow B$, which transforms a_i to the b_i , where $1 \leq i \leq k$ ".

For every word $w(x) \in W(X)$, where $X = \{x_1, \dots, x_k\}$ and every $C \in \Theta$ we can define a k -ary operation $w_C^*(c) = w(c)$ (in full notation $c = (c_1, \dots, c_k) \in C^k$, $x = (x_1, \dots, x_k) \in (W(X))^k$) or, more formal, $w_C^*(c) = \gamma_c(w(x))$, where γ_c is a well defined homomorphism $\gamma_c : W(X) \ni x \rightarrow c \in C$ (in full notation: homomorphism $\gamma_c : W(X) \rightarrow C$, which transforms x_i to the c_i for $1 \leq i \leq k$). This operation we call the **verbal operation** induced on the algebra C by the word $w(x) \in W(X)$. A little more detailed discussion about definition of the verbal operation you can see in [10, Section 2.1].

Now we will consider two kinds of substances, which will be also important for our method:

1. systems of bijections $\{s_B : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$ which fulfills these two conditions:

B1) for every homomorphism $\alpha : A \rightarrow B$ ($A, B \in \text{Ob}\Theta^0$) the mappings $s_B \alpha s_A^{-1}$ and $s_B^{-1} \alpha s_A$ is also a homomorphism;

B2) $s_B|_X = id_X$ for every $B = W(X) \in \text{Ob}\Theta^0$.

2. systems of words $\{w_\omega(x) \in W(X_\omega) \mid \omega \in \Omega\}$ which fulfills these two conditions:

Op1) $X_\omega = \{x_1, \dots, x_k\}$, where k is an arity of ω , for every $\omega \in \Omega$;

Op2) for every $B = W(X) \in \text{Ob}\Theta^0$ there exists an isomorphism $\sigma_B : B \rightarrow B^*$ (algebra B^* has the same domain as the algebra B and its operations ω_B^* are induced by $w_\omega(x)$ for every $\omega \in \Omega$) such as $\sigma_B|_X = id_X$.

The system of bijections $\{s_B^\Phi : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$, described in A2) and A3) of the definition of the strongly stable automorphism fulfills conditions B1) and B2) with $s_B = s_B^\Phi$.

We take a system of bijections $S = \{s_B : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$ which fulfills conditions B1) and B2). If arity of $\omega \in \Omega$ is k , we take $X_\omega = \{x_1, \dots, x_k\} \subset X_0$. $A_\omega = W(X_\omega)$ - free algebra in Θ . We have that $\omega(x) \in A_\omega$ ($x = (x_1, \dots, x_k)$) so there exists $w_\omega(x) \in A_\omega$ such that

$$w_\omega(x) = s_{A_\omega}(\omega(x)). \quad (1.2)$$

The system of words $\{w_\omega(x) \in A_\omega \mid \omega \in \Omega\}$ we denote $\mathfrak{W}(S)$. This system fulfills condition Op1) by our construction, condition Op2) with $\sigma_B = s_B$ ($B \in \text{Ob}\Theta^0$) by [9, Theorem 3]. For every $C \in \Theta$ we denote ω_C^* the verbal operation induced on the algebra C by $w_\omega(x)$. C^* will be the algebra, which has the same domain as the algebra C and its operations are $\{\omega_C^* \mid \omega \in \Omega\}$. By [10, Proposition 3.1] one can conclude that for every $C \in \Theta$ the algebra C^* belongs to Θ .

Contrariwise, if we have a system of words $W = \{w_\omega(x) \mid \omega \in \Omega\}$, which fulfills conditions Op1) and Op2), then the isomorphisms $\sigma_B : B \rightarrow B^*$ ($B \in \text{Ob}\Theta^0$) are bijections. For the system of bijections $\{\sigma_B : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$, which we denote $\mathfrak{S}(W)$, B1) fulfills by [5, 1.8, Lemma 8] and [10, Corollary from Proposition 3.2]; B2) fulfills by construction of these bijections.

If we have a system of bijections $S = \{s_B : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$ which fulfills conditions B1) and B2) then we can define an automorphism $\Phi(S) = \Phi$ of the category Θ^0 : Φ preserves all objects of the category Θ^0 and acts on its morphisms according to formula (1.1) with $s_B^\Phi = s_B$. Obviously Φ fulfills conditions A1) - A3) with $\{s_B^\Phi : B \rightarrow B \mid B \in \text{Ob}\Theta^0\} = S$, i.e. Φ is a strongly stable automorphism. Actually, the two different systems of bijections $S_1 = \{s_{1,B}^\Phi : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$ and $S_2 = \{s_{2,B}^\Phi : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$ can provide by formula (1.1) the same action on homomorphisms and, so, the same strongly stable automorphism of the category Θ^0 .

Proposition 1.1. *Every strongly stable automorphism Φ of the category Θ^0 can be obtained as $\Phi(\mathfrak{S}(W))$ where $W = \{w_\omega(x) \in A_\omega \mid \omega \in \Omega\}$ is a system of words which fulfills conditions Op1) and Op2).*

Proof. Let Φ be a strongly stable automorphism of the category Θ^0 . The system of bijections $S = \{s_B^\Phi : B \rightarrow B \mid B \in \text{Ob}\Theta^0\}$ fulfills conditions B1) and B2). Lets consider $\{w_\omega(x) \in A_\omega \mid \omega \in \Omega\} = \mathfrak{W}(S)$. By [10, Proposition 3.3], $\mathfrak{W}(S)$ fulfills conditions Op1) and Op2). By [10, Proposition 3.4], $\mathfrak{S}(\mathfrak{W}(S)) = S$, so $\Phi(\mathfrak{S}(\mathfrak{W}(S))) = \Phi(S)$. $\Phi(S)$ and Φ both preserve all objects of Θ^0 and act on the morphisms of Θ^0 by the same system of bijections S according to formula (1.1). Therefore they coincide. ■

So, if we describe the strongly stable automorphisms of the category Θ^0 , we must concentrate on finding out the systems of words, which fulfill conditions Op1) and Op2). However, in describing this, we must remember, that different systems of words which fulfill conditions Op1) and Op2) can provide us the same automorphism, because different systems of bijections can provide us the same automorphism.

This method we apply to the variety $\mathfrak{N}_d$ of the all nilpotent class no more than d groups. The variety $\mathfrak{A}_2$ of the abelian groups with the exponent no more than 2 is a subvariety of $\mathfrak{N}_d$ for every $d \in \mathbb{N}$. So, by the second theorem of Fudzivara [7, III.7.6] the variety $\mathfrak{N}_d$ is an IBN variety.

2 Applying the method.

The free i -generated group in $\mathfrak{N}_d$ is denoted as NF_i^d . The group signature is $\{1, -1, \cdot\}$, where 1 is a 0-ary operation of the taking of the unit, -1 is a unary operation of taking an inverse element and $\cdot$ is a binary operation of the multiplication. So, by our method, we must find out the systems of the words

$$\{w_1, w_{-1}, w.\} \quad (2.1)$$

such that $w_1 \in NF_0^d$, $w_{-1} \in NF_1^d$, $w. \in NF_2^d$ (condition Op1)) and condition Op2) fulfills for all $NF_i^d \in \text{Ob}\mathfrak{N}_d^0$ ($i \in \mathbb{N}$). But, as it will be clear above, we can consider only the word $w(x, y) \in NF_2^d$ (x, y are generators of NF_2^d) which fulfills this condition:

Op^d) If $(NF_2^d)^*$ is a set NF_2^d with the binary verbal operation induced by $w(x, y) \in NF_2^d$ (this operation we denote as " $*$ "), then in $(NF_2^d)^*$ are fulfilled all group axioms and exists an isomorphism $\sigma_d : NF_2^d \rightarrow (NF_2^d)^*$, such that $\sigma_d(x) = x$, $\sigma_d(y) = y$ (x, y are the generators of NF_2^d).

Proposition 2.1. *If in $(NF_2^d)^*$ are fulfilled all group axioms then $w(x, y) = xyg_2(x, y)$ ($g_2(x, y) \in \gamma_2(NF_2^d)$, $\gamma_i(G)$ is a i -th group in the lowest central series of the group G), $1_* = 1$, $a^{*k} = a^k$, for every $a \in NF_2^d$, $k \in \mathbb{Z}$ and $\gamma_i((NF_2^d)^*) \subset \gamma_i(NF_2^d)$ for every $i \in \mathbb{N}$ (1_* and a^{*k} is the unit of the $(NF_2^d)^*$ and degree of the element a according to the new operation).*

Proof. Every word in NF_2^d can be written as $w(x, y) = x^t y^s g_2(x, y)$ ($t, s \in \mathbb{Z}$). So we assume that $x * y = x^t y^s g_2(x, y)$. Then $1 * 1 = 1$, so $1 = 1_*$. Therefore

$x = x * 1 = x^t$, so $t = 1$. Analogously $s = 1$. For every $a \in NF_2^d$ and every $k_1, k_2 \in \mathbb{Z}$ we have $a^{k_1} * a^{k_2} = a^{k_1+k_2}$, because $g_2(a^{k_1}, a^{k_2})$ is calculated in the commutative group $\langle a \rangle$.

So, for every $a, b \in NF_2^d$ holds $(a, b)_* = a^{-1}b^{-1}g_2(a^{-1}, b^{-1}) * abg_2(a, b) = a^{-1}b^{-1}abl_2 = (a, b)l_2 \in \gamma_2(NF_2^d)$ ($l_2 \in \gamma_2(NF_2^d)$, (a, b) is a commutator of a and b and $(a, b)_*$ is a commutator of a and b according to the new operation $"*"$). We assume that for $k < i$ it is proved that $\gamma_k((NF_2^d)^*) \subset \gamma_k(NF_2^d)$. If $l_{i-1} \in \gamma_{i-1}((NF_2^d)^*)$, $b \in NF_2^d$, then we have

$$\begin{aligned} (l_{i-1}, b)_* &= l_{i-1}^{-1}b^{-1}g_2(l_{i-1}^{-1}, b^{-1}) * l_{i-1}bg_2(l_{i-1}, b) = \\ &= l_{i-1}^{-1}b^{-1}g_2(l_{i-1}^{-1}, b^{-1})l_{i-1}bg_2(l_{i-1}, b)g_2(l_{i-1}^{-1}b^{-1}g_2(l_{i-1}^{-1}, b^{-1}), l_{i-1}bg_2(l_{i-1}, b)) \cdot \\ &g_2(l_{i-1}^{-1}, b^{-1}), g_2(l_{i-1}, b) \in \gamma_i(NF_2^d), \text{ because } l_{i-1} \in \gamma_{i-1}(NF_2^d). \text{ So} \\ (l_{i-1}, b)_* &\equiv l_{i-1}^{-1}b^{-1}l_{i-1}bg_2(l_{i-1}^{-1}b^{-1}, l_{i-1}b) \pmod{\gamma_i(NF_2^d)}. \\ (l_{i-1}^{-1}b^{-1}, l_{i-1}b) &\equiv (l_{i-1}^{-1}, b)(b^{-1}, l_{i-1}) \equiv 1 \pmod{\gamma_i(NF_2^d)}. \end{aligned}$$

$g_2(x, y)$ has the weight 2 or more, so $g_2(l_{i-1}^{-1}b^{-1}, l_{i-1}b) \in \gamma_i(NF_2^d)$ and $(l_{i-1}, b)_* \in \gamma_i(NF_2^d)$. ■

Corollary 1. *If the system of words (2.1) fulfills conditions Op1) and Op2) then $w_1 = 1$, $w_{-1} = x^{-1}$.*

By this Corollary we can concentrate on a research of the verbal binary operations in the NF_2^d . We will find out the word $w(x, y) \in NF_2^d$, which fulfill condition Op^d).

Corollary 2. *In the condition of the Proposition 2.1 the group $(NF_2^d)^*$ is also a nilpotent class d group.*

By this Corollary, if in $(NF_2^d)^*$ are fulfilled all group axioms then the homomorphism $\sigma_d : NF_2^d \ni \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} x \\ y \end{pmatrix} \in (NF_2^d)^*$ is well defined.

Proposition 2.2. *The verbal operation induced on NF_2^2 by the word $w(x, y) = xy(y, x)^m$ fulfills the group axioms for every $m \in \mathbb{Z}$, but the homomorphism $\sigma_2 : NF_2^2 \rightarrow (NF_2^2)^*$ is an isomorphism if and only if $m = 0$ or $m = 1$, i.e., $w(x, y) = xy$ or $w(x, y) = yx$.*

Proof. Let $x * y = xy(y, x)^m$. We have $1_* = 1$, $a^{*(-1)} = a^{-1}$. For every $a, b, c \in NF_2^2$ we have

$$\begin{aligned} (a * b) * c &= ab(b, a)^m * c = \\ &= ab(b, a)^m c(c, ab(b, a))^m = abc(b, a)^m (c, a)^m (c, b)^m. \end{aligned}$$

$$\begin{aligned}
a * (b * c) &= a * bc(c, b)^m = \\
&= abc(c, b)^m (bc(c, b)^m, a)^m = abc(c, b)^m (b, a)^m (c, a)^m.
\end{aligned}$$

So $(a * b) * c = a * (b * c)$.

$$\begin{aligned}
(y, x)_* &= y^{-1} * x^{-1} * y * x = y^{-1} x^{-1} (x^{-1}, y^{-1})^m * yx(x, y)^m = \\
&= (y, x)^{1-2m} (yx, y^{-1} x^{-1})^m = (y, x)^{1-2m} ((y, x^{-1})(x, y^{-1}))^m = (y, x)^{1-2m}. \\
\sigma_2(x^{\alpha_1} y^{\alpha_2} (y, x)^{\alpha_3}) &= \sigma_2(x^{\alpha_1}) * \sigma_2(y^{\alpha_2}) * \sigma_2((y, x)^{\alpha_3}) = x^{\alpha_1} * y^{\alpha_2} * ((y, x)_*)^{\alpha_3} = \\
&= x^{\alpha_1} * y^{\alpha_2} * ((y, x)_*)^{\alpha_3} = x^{\alpha_1} * y^{\alpha_2} * (y, x)^{(1-2m)\alpha_3} = x^{\alpha_1} y^{\alpha_2} (y, x)^{\alpha_1 \alpha_2 m + (1-2m)\alpha_3} \\
&(\alpha_1, \alpha_2, \alpha_3 \in \mathbb{Z}). \text{ If } 1 - 2m \neq \pm 1, \text{ i.e., } m \neq 0, 1, \text{ then } (y, x) \notin \sigma_2(NF_2^d). \blacksquare
\end{aligned}$$

Proposition 2.3. *If $(NF_2^d)^*$ as in the condition Op^d , then $\gamma_i(NF_2^d) = \gamma_i((NF_2^d)^*)$ for every $i \in \mathbb{N}$.*

Proof. By Corollary 1 from Proposition 2.1, the operations "1" and "-1" of the group signature in NF_2^d and in $(NF_2^d)^*$ coincide. By [10, Proposition 3.2], original operations in NF_2^d are the verbal operations induced on NF_2^d by the words with respect to the operations "1", "-1" and "*". So we have a situation of Proposition 2.1 in which the group NF_2^d and NF_2^d changed they place. Hence, $\gamma_i(NF_2^d) \subset \gamma_i((NF_2^d)^*)$ for every $i \in \mathbb{N}$. $\blacksquare$

Lemma 2.1. *If the word $w(x, y)$ fulfills conditions Op^d , then the word $\kappa(w(x, y)) \in NF_2^{d-1}$ fulfills conditions Op^{d-1} ($\kappa : NF_2^d \rightarrow NF_2^{d-1}$ is the natural epimorphism).*

Proof. $NF_2^{d-1} \in \mathfrak{N}_d$. So on NF_2^{d-1} we can induce the operations of the group signature by words $1 \in NF_0^d$, $x^{-1} \in NF_1^d$, $w(x, y) \in NF_2^d$. This operations we denote by "1", "-1" and "o" correspondingly. NF_2^{d-1} with this operations we denote $(NF_2^{d-1})^\circ$.

$$\begin{array}{ccccccc}
1 & \rightarrow & \gamma_d(NF_2^d) & \hookrightarrow & NF_2^d & \xrightarrow{\kappa} & NF_2^{d-1} & \rightarrow & 1 \\
& & \downarrow \sigma_d & & \downarrow \sigma_d & & \downarrow \sigma_{d-1} & & \\
1 & \rightarrow & \gamma_d(NF_2^d) & \hookrightarrow & (NF_2^d)^* & \xrightarrow{\kappa} & (NF_2^{d-1})^\circ & \rightarrow & 1
\end{array} \tag{2.2}$$

The operations in $(NF_2^d)^*$ are induced by same words; $\kappa : NF_2^d \rightarrow NF_2^{d-1}$ is a homomorphism, so, by [5, 1.8, Lemma 8], κ is also a homomorphism from $(NF_2^d)^*$ to $(NF_2^{d-1})^\circ$. Therefore $(NF_2^{d-1})^\circ$ is an homomorphic image of the group $(NF_2^d)^*$, hence it is also a group. For every $a, b \in NF_2^{d-1}$ we have $a \circ b = w(a, b) = \alpha w(x, y)$, where $\alpha : NF_2^d \ni \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} a \\ b \end{pmatrix} \in NF_2^{d-1}$. We can also consider a homomorphism $\tilde{\alpha} : NF_2^{d-1} \ni \begin{pmatrix} \kappa(x) \\ \kappa(y) \end{pmatrix} \rightarrow \begin{pmatrix} a \\ b \end{pmatrix} \in NF_2^{d-1}$.

We have $\alpha = \tilde{\alpha}\kappa$, so $a \circ b = \tilde{\alpha}(\kappa w(x, y))$. Therefore the operation "o" is induced in NF_2^{d-1} by the word $\kappa w = w(\kappa(x), \kappa(y)) \in NF_2^{d-1}$ ($\kappa(x)$ and $\kappa(y)$ are free generators of NF_2^{d-1}). This operation fulfills all group axioms, hence, by Corollary 2 from Proposition 2.1, $(NF_2^{d-1})^\circ$ is also a nilpotent class $d-1$ group. So $\sigma_{d-1} : NF_2^{d-1} \ni \begin{pmatrix} \kappa(x) \\ \kappa(y) \end{pmatrix} \rightarrow \begin{pmatrix} \kappa(x) \\ \kappa(y) \end{pmatrix} \in (NF_2^{d-1})^\circ$ is a well defined homomorphism. Our goal is to prove that it is an isomorphism.

By Proposition 2.3, we have $\sigma_d(\gamma_d(NF_2^d)) \subset \gamma_d((NF_2^d)^*) = \gamma_d(NF_2^d)$. By consideration of the acting of σ_d^{-1} , we achieve $\gamma_d(NF_2^d) \subset \sigma_d(\gamma_d(NF_2^d))$. In the diagram (2.2) we have two exact rows, because $\ker \kappa = \gamma_d(NF_2^d)$. The left square of this diagram is commutative. The right square is commutative too, because σ_d and σ_{d-1} preserve the generators of the corresponding free nilpotent groups. σ_d is an isomorphism, so σ_{d-1} too. ■

Lemma 2.2. *For every $d \geq 2$ there are only two words: $w = xy$ and $w = yx$ from NF_2^d which fulfills condition Op^d .*

Proof. For $d = 2$ it is proved in the Proposition 2.2. We will assume that it is proved for all natural numbers lesser then d .

Let $w(x, y) \in NF_2^d$ fulfills condition Op^d and $*$ be an operation in NF_2^d induced by $w(x, y)$. By Lemma 2.1 and assumption of induction $\kappa w(x, y) = \kappa(x)\kappa(y)$ or $\kappa w(x, y) = \kappa(y)\kappa(x)$.

In the first case we have $w(x, y) = xyr(x, y)$, where $r(x, y) \in \gamma_d(NF_2^d)$. For every $a, b, c \in NF_2^n$ we have $(a * b) * c = abr(a, b) * c = abr(a, b)cr(abr(a, b), c) = abcr(a, b)r(ab, c)$ ($r(a, b) = \alpha(r(x, y))$ where $\alpha : NF_2^d \ni \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} a \\ b \end{pmatrix} \in NF_2^d$ for every $a, b \in NF_2^d$, so $r(a, b) \in \gamma_d(NF_2^d) \subset Z(NF_2^d)$). Also we have $a * (b * c) = a * bcr(b, c) = abcr(b, c)r(a, bcr(b, c)) = abcr(b, c)r(a, bc)$. $*$ fulfills the axiom of associativity, so

$$r(a, b)r(ab, c) = r(b, c)r(a, bc). \quad (2.3)$$

We will prove that for $d \geq 3$ there isn't any nontrivial word $r(x, y)$, which is generated by commutators of the weight d and fulfills the condition (2.3). We will shift all our calculation from the free nilpotent group NF_2^d to the NL_2^d - free nilpotent class n Lie algebra over $\mathbb{Q}$ with generators x and y . By [1, 8.3.9], $\sqrt{NF_2^d} = (NL_2^d)^\circ$, where $\sqrt{NF_2^d}$ is a Maltsev completion of the group NF_2^d and $(NL_2^d)^\circ$ is a group which coincides with NL_2^d as a set and has a multiplication defined by Campbell-Hausdorff formula. By consideration of the Campbell-Hausdorff formula we have that $r(a, b) = q(a, b)$, where $q(x, y)$ is the word in NL_2^d , which we achieve from $r(x, y)$ by replacement of the circular brackets of the group commutators by Lie brackets and multiplication in NF_2^d by addition in NL_2^d , $q(a, b)$ is a result of substitution of the a and b instead x and y correspondingly ($a, b \in NF_2^d$) to the word $q(x, y)$. Also by consideration of the Campbell-Hausdorff formula $ab \equiv a + b \pmod{(NL_2^d)^2}$ for every $a, b \in$

NF_2^d ($(NL_2^d)^i$ is the i -th ideal of the lowest central series of the algebra NL_2^d , $1 \leq i \leq d$) and $ab = a + b$ if $a, b \in \gamma_d(NF_2^d)$. Therefore the condition (2.3) we can write as

$$q(a, b) + q(a + b, c) = q(b, c) + q(a, b + c). \quad (2.4)$$

We shall substitute $a = \lambda x$, $b = x$, $c = y - (\lambda \in \mathbb{Z})$ to (2.4) and we will achieve:

$$q((\lambda + 1)x, y) = q(x, y) + q(\lambda x, x + y). \quad (2.5)$$

NL_2^d is a direct sum of its polyhomogeneous (homogeneous as according x , as according y separately) components (proof of this fact is similar to the proof of [1, 2.2.5]). Let $q(x, y) = \sum_{i=1}^{d-1} q_i(x, y)$, where $q_i(x, y)$ is a polyhomogeneous component of $q(x, y)$ corresponding to the degree i of x and $d - i$ of y . By (2.5) we have

$$\sum_{i=1}^{d-1} (\lambda + 1)^i q_i(x, y) = \sum_{i=1}^{d-1} q_i(x, y) + \sum_{i=1}^{d-1} \lambda^i q_i(x, x + y). \quad (2.6)$$

When we develop $q_i(x, x + y)$ by additivity, we achieve $q_i(x, x + y) = q_i(x, y) + \sum_{j=i+1}^{d-1} m_{i,j}(x, y)$ where $m_{i,j}(x, y)$ is a polyhomogeneous component of $q_i(x, x + y)$ with the degree j of x and $d - j$ of y . In particular $q_1(x, x + y) = q_1(x, y) + \sum_{j=2}^{d-1} m_{1,j}(x, y)$. By comparison of the polyhomogeneous components of (2.6) with the degree 2 of x and $d - 2$ of y we have

$$\left((\lambda + 1)^2 - 1 - \lambda^2 \right) q_2(x, y) = \lambda^2 m_{1,2}(x, y). \quad (2.7)$$

Let $\{e_1(x, y), \dots, e_k(x, y)\}$ be a linear basis of the d -th ideal of the lowest central series of NL_2^d and $q_2(x, y) = \sum_{s=1}^k \rho_s e_s(x, y)$, $m_{1,2}(x, y) = \sum_{s=1}^k \mu_s e_s(x, y)$ ($\rho_s, \mu_s \in \mathbb{Z}$). From (2.7) we have

$$\left((\lambda + 1)^2 - 1 - \lambda^2 \right) \rho_s = \lambda^2 \mu_s$$

($s \in \{1, \dots, k\}$). We take two values of λ such that we will achieve $\rho_s = 0$, $\mu_s = 0$. Hence $q_2(x, y) = 0$, $q_2(x, x + y) = 0$, $m_{1,2}(x, y) = 0$. Step by step, analogously we conclude, that $q_i(x, y) = 0$, $q_i(x, x + y) = 0$, $m_{1,i}(x, y) = 0$ for $i \in \{3, \dots, d - 1\}$. Therefore $q(x, y) = q_1(x, y)$. Now we shall substitute $a = x$, $b = y$, $c = \lambda y - (\lambda \in \mathbb{Z})$ to (2.4) and we will achieve:

$$q_1(x, y) + q_1(x + y, \lambda y) = q_1(x, (\lambda + 1)y)$$

or

$$\left((\lambda + 1)^{d-1} - \lambda^{d-1} - 1 \right) q_1(x, y) = 0,$$

so $q_1(x, y) = 0$, $q(x, y) = 0$, $r(x, y) = 1$ and $w(x, y) = xy$.

Analogously we consider the case $w(x, y) = yx$. ■

Theorem 2.1. *All automorphisms of the category of the nilpotent class d free groups are inner.*

Proof. Let $W = \{w_1, w_{-1}, w.\}$ be the system of words, which fulfills conditions Op1) and Op2). By Lemma 2.2 there are no more than two opportunities: $W = \{1, x^{-1}, xy\}$ or may also be $W = \{1, x^{-1}, yx\}$. It is easy to check that both these systems actually fulfill conditions Op1) and Op2). By Proposition 1.1, all strongly stable automorphisms of $\mathfrak{N}_d^0$ can be achieved as $\Phi(\mathfrak{S}(W))$. If W fulfills conditions Op1) and Op2), then $\mathfrak{S}(W)$ fulfills conditions B1) and B2) (see Section 1), so, by [9, Lemma 3 and Theorem 2], $\Phi(\mathfrak{S}(W))$ is an inner automorphism if and only if for every $B \in \text{Ob}\mathfrak{N}_d^0$ exists an isomorphism c_B from B to B^* , where B^* is an algebra which has the same domain as algebra B and operations induced by the $\mathfrak{W}(\mathfrak{S}(W)) = W$ (see [10, Proposition 3.4]), such that $c_B \alpha c_B^{-1} = \alpha$ for every homomorphism $\alpha : A \rightarrow B$ ($A, B \in \text{Ob}\Theta^0$). If $W = \{1, x^{-1}, xy\}$, then $B^* = B$ and we can take $c_B = id_B$ for every $B \in \text{Ob}\mathfrak{N}_d^0$. And if $W = \{1, x^{-1}, yx\}$, then for every $B \in \text{Ob}\mathfrak{N}_d^0$ we can take $c_B : B \rightarrow B^*$ such that $c_B(b) = b^{-1}$ for every $b \in B$. ■

3 Acknowledgements.

I dedicate this paper to the 80th birthday of Prof. B. Plotkin. He motivated this research and was very heedful to it. Of course, I must appreciate to Prof. B. Plotkin and Dr. G. Zhitomirski for their marvelous paper [9]. This research could not take place without their deep results. Prof. S. Margolis also was very heedful to my research. Very useful discussion with Prof. L. Rowen, Dr. R. Lipyanski, Dr. E. Plotkin and Dr. G. Zhitomirski, also help me in the writing of this paper.

References

- [1] Ju. Bahturin, *Identities in Lie algebras*, (Russian), Nauka, Moscow, 1985.
- [2] A. Berzins, B. Plotkin, E. Plotkin, *Algebraic geometry in varieties of algebras with the given algebra of constants*, Journal of Math. Sciences, **102:3**, (2000), pp. 4039 – 4070.
- [3] J. Dyer, E. Formanek, *The automorphism group of a free group is complete*, J. London Math. Soc. (2), **11:2** (1975), pp. 181 – 190.
- [4] E. Formanek, *A question of B. Plotkin about semigroup of endomorphisms of a free group*, Proc. Amer. Math. Soc., **130** (2002), pp. 935 – 937.
- [5] G. Grätzer, *Universal algebra*. Second edition. Springer-Verlag, 1979.
- [6] M. Kassabov, *On the Automorphism Tower of Free Nilpotent Groups*, <http://arxiv.org/abs/math.GR/0311488>.

- [7] A. Kurosh, *Lectures in general algebra* (Russian), Fizmatgiz, Moscow, 1962.
- [8] Plotkin B. *Algebras with the same (algebraic) geometry*, Proceedings of the International Conference on Mathematical Logic, Algebra and Set Theory, dedicated to 100 anniversary of P.S.Novikov, Proceedings of the Steklov Institute of Mathematics, MIAN, v.242, (2003), pp. 17 – 207.
- [9] B. Plotkin, G. Zhitomirski, *On automorphisms of categories of free algebras of some varieties*, <http://arxiv.org/abs/math.RA/0501331>.
- [10] A. Tsurkov *Automorphic equivalence of one-sorted algebras*, <http://arxiv.org/abs/math.GM/0509032>.